Author index

Abdel-Sattar, M.A., see Shaw *et al.*, 552

Aharonson, N., see Yarden *et al.*, 69

Ahl, Patricia, Gianinazzi, S., Samson, Regine and Benjama, A. Cultivar dependence of polyacrylic acid effects on *Pseudomonas syringae* in *Nicotiana tabacum*, 221

Alivizates, A.S. Bacterial wilt of tomato in Greece caused by *Erwinia chrysanthemi*, 638

Allen, E.C., see Jordan, 200

Anthony, Vivienne M., Shattock, R.C. and Williamson, B. Life-history of *Phragmidium rubi-idaei* on red raspberry in the United Kingdom, 510

Anthony, Vivienne M., Shattock, R.C. and Williamson, B. Interaction of red raspberry cultivars with isolates of *Phragmidium rubi-idaei*, 521

Asman, Ariful, see Bennett *et al.*, 487; see Hunt *et al.*, 495

Aubury, R.G., see Lockwood *et al.*, 341; see Jellis *et al.*, 347

Bainbridge, A., Fitt, B.D.L., Creighton, N.F. and Cayley, G.R. Use of fungicides to control chocolate spot (*Botrytis fabae*) on winter field beans (*Vicia faba*), 5

Ball, Elizabeth, see Gay *et al.*, 353

Bambridge, Judith M., see Maude, 435

Barbetti, Martin J. Survey of fungi associated with subterranean clover leaves and petioles in Western Australia, 49

Bastos, C.N. and Evans, H.C. A new pathotype of *Crinipellis perniciosa* (witches' broom disease) on solanaceous hosts, 306

Bautista, Rosemary M., see Imperial *et al.*, 391

Benedikz, P.W., see Scott *et al.*, 578

Benjama, A., see Ahl *et al.*, 221

Bennett, Chris P.A., Hunt, Peter and Asman, Ariful. Association of a xylem-limited bacterium with Sumatra disease of cloves in Indonesia, 481

Bennett, Chris P.A., see Hunt *et al.*, 495

Best, G.R., see Jordan, 200

Blunt, Sarah, see Hill, 472

Boag, B. and Topham, P.B. The use of associations of nematode species to aid the detection of small numbers of virus-vector nematodes, 20

Brunt, A.A., see Phillips, Sue, 440

Byford, W.J. A comparison of fungicide seed treatments to improve sugar beet seedling establishment, 463

Caten, C.E., see Newton, 235

Cayley, G.R., see Bainbridge *et al.*, 5

Chase, A.R. Bacterial leaf spot of *Codiaeum variegatum* cultivars caused by *Xanthomonas campestris* pv. *poinsettiicola*, 446

Chet, I., see Strashnow *et al.*, 146

Clarkson, John M. and Heale, J.B. Pathogenicity and colonization studies on wild-type and auxotrophic isolates of *Verticillium albo-atrum* from hop, 119

Clarkson, John M. and Heale, J.B. Heterokaryon compatibility and genetic recombination within a host plant between hop wilt isolates of *Verticillium albo-atrum*, 129

Cole, A.L.J., see Cromey, 83

Coley-Smith, J.R. Methods for the production and use of sclerotia of *Sclerotium cepivorum* in field germination studies, 380

Cooke, R.C., see Lutchmeah, R.S., 528

Cooper, J.I., see Edwards *et al.*, 95

Cornford, C.A. and Pitt, D. The nature of virulence in isolates of *Bremia lactucae* from South Devon, 38

Creighton, N.F., see Bainbridge *et al.*, 5

Cromey, M.G. & Cole, A.L.J. Cytology of the host-pathogen interactions between *Lolium perenne* and *Drechslera dictyoides*, 83

Crosse, J.E., see Kairu *et al.*, 207

Crute, I.R., see Norwood, 481

Davies, R.M. and Heale, J.B. *Botrytis cinerea* in stored cabbage: the use of germ-tube growth on leaf discs as an indicator of potential head rot, 408

Della Vecchia, P.T., see Reifschneider, 300

Edwards, M.-L., Cooper, J.I., Massalski, P.R. and Green, B. Some properties of a virus-like agent found in *Brachypodium sylvaticum* in the United Kingdom, 95

Elad, Y., see Strashnow *et al.*, 146

Evans, H.C., see Bastos, 306

Fitt, B.D.L., see Bainbridge *et al.*, 5

Fletcher, J.T. and Lord, Karen L. A stem rot of tomato caused by *Fusarium merismoides*, 443

Flett, J.D., see Wimalajeewa, 248

Ford, M.A., see Scott *et al.*, 578

Francki, R.I.B. and Miles, R. Mechanical transmission of sowbane mosaic virus carried on pollen from infected plants, 11

Fyfe, A.M., see Shaw *et al.*, 552

Garrett, R.G. see Moran *et al.*, 467

Gay, John L., Martin, Marion and Ball, Elizabeth. The impermeability of powdery mildew conidia and their germination in arid environments, 353

Gianinazzi, S., See Ahl *et al.*, 221

Gibbs, A.J., see Guy, 532

Gill, P.A. Lateral bud necrosis in the black currant (*Ribes nigrum* L.), 297

da Graça, J.V., Trench, T.N. and Martin, M.M. Tomato spotted wilt virus in commercial Cape gooseberry (*Physalis peruviana*) in Transkei, 451

Green, B., see Edwards *et al.*, 95

Gregory, S.C. The use of potato tubers in pathogenicity studies of *Armillaria* isolates, 41

Griffin, M.J., see King, 272

Guy, P.L. and Gibbs, A.J. Further studies on turnip yellow mosaic tymovirus isolates from an endemic Australian *Cardamine*, 532

Hall, K. and Kavanagh, J.A. Light and scanning electron microscope studies of leaf blotch of onion caused by *Cladosporium alii-cepae*, 1

Heale, J.B., see Clarkson, 119, 129; see Davies, 408

Hammond, Kim E., Lewis, B.G. and Musa, T.M. A systemic pathway in the infection of oilseed rape plants by *Leptosphaeria maculans*, 557

Hibberd, P.G., see Shaw *et al.*, 552

Hill, S.A. and Blunt, Sarah. Detection of two viruses infecting *Narcissus* by enzyme-linked immunosorbent assay, 472

Hoffmann, R.M., see Turner, 54

Hollins, T.W., Scott, P.R. and Paine, J.R. Morphology, benomyl resistance and pathogenicity to wheat and rye of isolates of *Pseudocercosporella herpotrichoides*, 379

Holmes, S.J.I. Barley yellow dwarf virus in ryegrass and its detection by ELISA, 214

Hughes, Fiona L. and Rijkenberg, F.H.J. Scanning electron microscopy of early infection in the uredial stage of *Puccinia sorghi* in *Zea mays*, 61

Humpherson-Jones, F.M. The incidence of *Alternaria* spp. and *Leptosphaeria maculans* in commercial brassica seed in the United Kingdom, 385

Hunt, Peter, see Bennett *et al.*, 487

Hunt, Peter, Bennett, Chris P.A. and Asman, Ariful. Suppression of Sumatra disease symptoms in cloves treated with antibiotics, 495

Imperial, Julita S., Bautista, Rosemary M. and Randles, John W. Transmission of the coconut cadang-cadang viroid to six species of palm by inoculation with nucleic acid extracts, 391

Ioannou, N. Yellow leaf curl and other virus diseases of tomato in Cyprus, 428

Ishii, Hideo, Udagawa, Hideo, Yanase, Haruo and Yamaguchi, Akira. Resistance of *Venturia nashicola* to thiophanate-methyl and benomyl: build-up and decline of resistance in the field, 363

Jellis, G.J., see Lockwood *et al.*, 341

Jellis, G.J., Lockwood, G. and Aubury, R.G. Phenotypic influences on the incidence of infection by *Ascochyta fabae* in spring varieties of faba beans (*Vicia faba*), 347

Johnson, R., see Newton *et al.*, 225

Jones, H.G., see Scott *et al.*, 578

Jones, R.A.C. Further studies on resistance-breaking strains of potato virus X, 182

Jordan, V.W.L., Best, G.R. and Allen, E.C. Effects of *Pyrenophora teres* on dry matter production and yield components of winter barley, 200

Kairu, G.M., Nyangena, C.M.S. and Crosse, J.E. The effect of copper sprays on bacterial blight and coffee berry disease in Kenya, 207

Katan, J., see Yarden *et al.*, 69

Kavanagh, J.A., see Hall, 1

King, J.E. and Griffin, M.J. Survey of benomyl resistance in *Pseudocercosporella herpotrichoides* on winter wheat and barley in England and Wales in 1983, 272

Knight, V.H. A new indicator for gooseberry vein banding virus, 629

Koffman, W., see Penrose *et al.*, 228

Lam, A. Effect of fungal pathogens on digestibility and chemical composition of Italian ryegrass (*Lolium multiflorum*) and tall fescue (*Festuca multiflorum*), 190

Lapwood, D.H. and Read, P.J. A simplified slice method for assessing tuber susceptibility of potato cultivars to *Erwinia carotovora* subsp. *atroseptica*, 284

Law, C.N., see Pink, 255

Lewis, B.G., see Hammond *et al.*, 557

Lewis, J.A. and Papavizas, G.C. Characteristics of alginate pellets formulated with *Trichoderma* and *Gliocladium* and their effect on the proliferation of the fungi in soil, 571

Lewis, Susan, see MacNish, 159

Lockwood, G., Jellis, G.J. and Aubury, R.G. Genotypic influences on the incidence of infection by *Ascochyta fabae* in winter-hardy faba beans (*Vicia faba*), 341

Lockwood, G., see Jellis *et al.*, 347

Lord, Karen L., see Fletcher, 443

Lutchmeah, R.S. and Cooke, R.C. Pelleting of seed with the antagonist *Pythium oligandrum* for biological control of damping-off, 528

Madhosingh, Clarence and Orr, Winston. Zearalenone production in *Fusarium culmorum* after transformation with DNA of *F. graminearum*, 402

Malathrakis, N.E. Tomato crown and root rot caused by *Fusarium oxysporum* f. sp. *radicis-lycopersici* in Greece, 438

Manners, J.G., see McGregor, 263; see Osman-Ghani, 75

Mansfield, J.W., see Stewart, 25

MacNish, G.C. Mapping rhizoctonia patch in consecutive cereal crops in Western Australia, 165

MacNish, G.C. Methods of reducing rhizoctonia patch of cereals in Western Australia, 175

MacNish, G.C. and Lewis, Susan. Methods of measuring rhizoctonia patch of cereals in Western Australia, 159

McGregor, A.J. and Manners, J.G. The effect of temperature and light intensity on growth and sporulation of *Puccinia striiformis* on wheat, 263

Martin, Marion, see Gay *et al.*, 353

Martin, M.M., see da Graça *et al.*, 451

Massalski, P.R., see Edwards *et al.*, 95

Maude, R.B. and Bambridge, Judith M. Effects of seed treatments and storage on the incidence of *Phoma betae* and the viability of infected red beet seeds, 435

Merriman, P.R., see Porter, 108

Miles, R., see Francki, 11

Moran, Jane R., Wilson, Jenny M., Garrett, R.G. and Smith, P.R. ELISA indexing of commercial carnations for carnation mottle virus using a urease-antibody conjugate, 467

Musa, T.M., see Hammond *et al.*, 557

Nair, N.G. and Parker, Frances E. Midseason bunch rot of grapes: an unusual disease phenomenon in the Hunter Valley, Australia, 302

Nathaniels, N.Q.R. *Phoma valerianellae* on corn salad, 449

Newman, Pamela L. Variation amongst isozymes of *Rhynchosporium secalis*, 329

Newman, Pamela L. and Owen, H. Evidence of asexual recombination in *Rhynchosporium secalis*, 338

Newton, A.C., Caten, C.E. and Johnson, R. Variation for isozymes and double-stranded RNA among isolates of *Puccinia striiformis* and two other cereal rusts, 235

Nicholls, M.R., see Penrose *et al.*, 228

Nieves-Brun, Carmen. Infection of roots of *Dieffenbachia maculata* by the foliar blight and soft rot pathogen, *Erwinia chrysanthemi*, 139

Norwood, Judith M. and Crute, I.R. Further characteristics of field resistance in lettuce to *Bremia lactucae* (downy mildew), 481

Nyangena, C.M.S., see Kairu, 207

O'Neill, T.M. *Mycocentrospora acerina* causing leaf spot and dieback of anemone seedlings, 632

Orr, Winson, see Madhosingh, 402

Osman-Ghani, N. and Manners, J.G. Components of partial resistance to *Puccinia striiformis* in barley, 75

Owen, H., see Newman, 338

Paine, J.R., see Hollins *et al.*, 369

Papavizas, G.C., see Lewis, 571

Pappas, A.C. Metalaxyl resistance in *Phytophthora infestans* on greenhouse tomatoes in Greece, 293

Parker, Frances E., see Nair, 302

Pearce, R.B., see Woodward, 477

Penrose, L.J., Koffman, W. and Nicholls, M.R. Field occurrence of vinclozolin resistance in *Monilinia fructicola*, 228

Phelps, Kathleen, see Walkey *et al.*, 545

Phillips, Sue and Brunt, A.A. Occurrence of cucumber mosaic virus and asparagus virus II in asparagus (*Asparagus officinalis* L. var. *officinalis*) in Britain, 440

Pink, D.A.C. and Law, C.N. The effect of homoeologous group 7 chromosomes upon adult plant resistance of wheat to yellow rust (*Puccinia striiformis*), 255

Pitt, D., see Cornford, 38

Polley, R.W. The relationship between brown and corky root rot caused by *Pyrenochaeta lycopersici* and yield loss in tomatoes established by the use of artificially infested soils, 502

Porter, I.J. and Merriman, P.R. Evaluation of soil solarization for control of root diseases of row crops in Victoria, 108

Presly, A.H. Studies on *Botrytis* spp. occurring on onions (*Allium cepa*) and leeks (*Allium porrum*), 422

Prior, C. Cocoa quarantine: measures to prevent the spread of vascular-streak dieback in planting material, 603

Randles, John W., see Imperial *et al.*, 391

Read, P.J., see Lapwood, 284

Reifschneider, F.J.B. and Della Vecchia, P.T. Thiabendazole as an iatrogenic agent in carrot black rot, 300

Rijkenberg, F.H.J., see Hughes, 61

Rocha, H.M. and Wheeler, B.E.J. Factors influencing the production of basidiocarps and the deposition and germination of basidiospores of *Crinipellis perniciosa*, the causal fungus of witches' broom on cocoa (*Theobroma cacao*), 319

Rudgard, S.A. and Wheeler, B.E.J. Relationships between establishment of *Erysiphe cruciferarum* and soluble amino-acids in leaves of Brussels sprouts, 609

Rudgard, S.A. and Wheeler, B.E.J. The development of *Erysiphe cruciferarum* on field-grown Brussels sprouts and associated changes in soluble amino-acids in foliage leaves, 616

Samson, Regine, see Ahl *et al.*, 221

Scott, E.S. Production and characterization of single-spore isolates of *Plasmodiophora brassicae*, 287

Scott, P.R., Benedikz, H.G., Jones, H.G. and Ford, M.A. Some effects of canopy structure and microclimate on infection of tall and short wheats by *Septoria nodorum*, 578

Scott, P.R., see Hollins *et al.*, 369

Shattock, R.C., see Anthony *et al.*, 510, 521

Shaw, Charles G. III. *In vitro* responses of different *Armillaria* taxa to gallic acid, tannic acid and ethanol, 594

Shaw, D.S., Fyfe, A.M., Hibberd, P.G. and Abdal-Sattar, M.A. Occurrence of the rare A2 mating type of *Phytophthora infestans* on imported Egyptian potatoes and the production of sexual progeny with A1 mating types from the UK, 552

Sijaona, M.E.R., see Teri, 636

Sivan, A., see Strashnow *et al.*, 146

Smith, P.R., see Moran *et al.*, 467

Stewart, Alison and Mansfield, J.W. The composition of wall alterations and appositions (reaction

material) and their role in the resistance of onion bulb scale epidermis to colonization by *Botrytis allii*, 25

Strashnow, Y., Elad, Y., Sivan, A. and Chet, I. Integrated control of *Rhizoctonia solani* by methyl bromide and *Trichoderma harzianum*, 146

Sutton, M.W. and Wale, S.J. The control of *Penicillium corymbiferum* on crocus and its effect on corm production, 566

Teri, J.M. and Sijaona, M.E.R. Tannia leaf spot in Tanzania, 636

Topham, P.B., see Boag, 20

Tosi, L., see Zazzerini, 415

Trench, T.N., see da Graça *et al.*, 451

Turner, J.G. and Hoffman, R.M. Effect of the polygalacturonase inhibitor from pea on the hydrolysis of pea cell walls by the endopolygalacturonase from *Ascochyta pisi*, 54

Udagawa, Hideo, see Ishii *et al.*, 363

Wale, S.J., see Sutton, 566

Walkey, D.G.A., Ward, C.M. and Phelps, Kathleen. Studies on lettuce mosaic virus resistance in commercial lettuce cultivars, 545

Ward, C.M., see Walkey *et al.*, 545

Wheeler, B.E.J., see Rocha, 319

Wheeler, B.E.J., see Rudgard, 609; 616

Whitehead, A.G. The potential value of British wild *Solanum* spp. as trap crops for potato cyst-nematodes, *Globodera rostochiensis* and *G. pallida*, 105

Williamson, B., see Anthony *et al.*, 510; 521

Wilson, Jenny M., see Moran *et al.*, 467

Wimalajeewa, D.L.S. and Flett, J.D. A study of populations of *Pseudomonas syringae* pv. *syringae* on stonefruits in Victoria, 248

Woodford, J.A.T. Cutworm damage to raspberry and black currant in Scotland, 626

Woodward, S. and Pearce, R.B. Methods for visualizing hyaline fungi for the detection of antifungal compounds on thin-layer chromatograms, 477

Yamaguchi, Akira, see Ishii *et al.*, 363

Yanasa, Haruro, see Ishii *et al.*, 363

Yarden, O., Katan, J. and Aharonson, N. A rapid bioassay for the determination of carbendazim residues in soil, 69

Zazzerini, A. and Tosi, L. Antagonistic activity of fungi isolated from sclerotia of *Sclerotinia sclerotiorum*, 415